## 75 mm

TMAKKA



## Sildenafil 100mg Oral Jelly

Each 5gm sachet contains: Sildenafil Citrate Eq. to Sildenafil......100 mg Approved colour used

## 4 pouches

Direction for Use : One sachet for one-time-use only

Sildenafil 100mg Oral Jelly ULTIMATE STAYING POWER



Registered Office 321777 State of Kuwait, Sharq, Block 5, Ahmad Al Jaber Street, Hamad Tower, Floor 12

BARCODE

## SACHET ORANGE Size: 70x50 mm

## ості пости пости

## FRONT

Sildenafil 100mg Oral Jelly Each 5gm sachet contains: Sildenafil Citrate Eq. to Sildenafil......00 mg Excipients......QS Approved colour used

## ULTIMATE STAYING POWER Quicker Onset of Action as compared to conventional Tablets

Diractions for uses • Open sachet and consume entire contents • One sachet for one lime use only • Use strictly under medical supervision • In Two Flavours (Pinsapple, Orange) • Dosage: As directed by the physician • Stare in a cool. dry & dark place

# 

SACHET PINEAPPLE

Size: 70x50 mm

## FRONT

BACK

Sildenafil 100mg Oral Jelly Each 5gm sachet contains: Sildenafil Citrate Eq. to Sildenafil......00 mg Excipients.....QS Approved colour used

ULTIMATE STAYING POWER Quicker Onset of Action as compared to conventional Tablets

### Directions for uses

 Open sachet and consume entire contents One sachet for one lime use only Use strictly under medical supervision In Two Flavours (Pineapple, Orange) Dosage: As directed by the physician Stare in a cool, dry & dark place

## BACK

Mfg. Lic. No: KD/561 Batch No.: Mfg. Date: Exp. Date:

Manufactured by: Maxwell Life Science Pvt. Ltd. T-53 & T-81, MIDC, TARAPUR, BOISAR District: Palghar - 401506 .

Manufactured for: CASTILLO INTERNATIONAL GROUP Registered Office 321777 State of Kuwait, Sharq, Block 5, Ahmad Al Jaber Street, Hamad Tower, Floor 12 تمخن TMAKKAN

Mfg. Lic. No: KD/561 Batch No.: Mfg. Date: Exp. Date:

Manufactured by: Maxwell Life Science Pvt. Ltd. T-53 & T-81, MIDC, TARAPUR, BOISAR District: Palghar - 401506 .

Manufactured for: CASTILLO INTERNATIONAL GROUP Registered Office 321777 State of Kuwait, Sharq, Block 5, Ahmad Al Jaber Street, Hamad Tower, Floor 12

## 250 mm x 140 mm FRONT BACK



have ever had heart problems (for example, angina, chest pain, heart failure, irregular heartbeat, heart attack, or narrowing of the aortic valve) have ever had a stroke have low or high blood pressure

have ever had severe vision loss

have a rare inherited eve disease called retinitis pigmentosa

have ever had kidney problems

Excipients.

have ever had liver problems

have ever had liver problems have ever had blood problems, including sickle cell anemia or leukemia are allergic to sildenafil (the active ingredient) have a deformed penis, Peronei's disease, or ever had an erection that lasted longer than 4 hour have stomach ulers or any kind of bleeding problems you are taking other medicines Some medicines can affect the way Seedios Oral Jelly works. Tell your doctor about the medications you are taking. Do not start or stop taking medicine before consulting your doctor. Seedios 0ral Jelly should never be used with medicines containing nitrates. If you take medicines called alpha-blockers for the treatment of high blood pressure or prostate problems, your

blood pressure may drop suddenly. You may feel dizzy or pass out.

Tournary technizes on high technizes on the adjusted. Seedios oral Jelly should not be used with other medical treatments that cause erections. These treatments include pills, drugs injected or inserted into the penis, implants or vacuum pumps. Seedios oral Jelly only for patients with arectife dysfunction. It is not for newborns, children or women. Seedios 0ral Jelly does not protect you or your partner against sexually transmitted diseases, including HIV - the virus that causes AIDS. It is not a hormone or an aphrodisiae.

## PREGNANCY, BREASTFEEDING AND FERTILITY:

Section on 1 below of the section of

## SIDE EFFECTS

SIDE EFFECTS: Most Common Effects: In pre-approval clinical trials with sildenafil, as detailed by eMedTV, the most common side effect was headache, experienced by up to 16% of participants in some studies. Other adverse reactions reported in pre-approval clinical trials and anecdotally post-approval include dizziness, facial flushing, indigestion, upset stomach, diarrhea, nasal congestion, and infection of the throats. urinary tract. Cardiovascular problems: Sildenafil can cause problems for people with cardiovascular disease, as noted in the Physicians Desktop. Reference (PDR). It has been associated with a fast or irregular heartheat and therefore may not be safe for people with theart disease, angina pectoris, low or high blood pressure, arrhythmia or anyone who has suffered a heart attack or stroke in the past six months. Vision side effects: Some men have experienced studen vision loss after taking sildenafil, according to the J.S. National Institutes of Health (NIH). This may involve a decrease in vision, or a partial or total loss of vision. The condition has been permanent in some cases. Other visual side effects reported by sildenafil users include blurred vision, eyes that are more sensitive to light, and changes in color vision, such as seeing a blue tint on objects or having difficulty divention blue of the from blue.

difficulty distinguishing green from blue.

Priapism: Sildenafil has been linked to priapism, a condition involving a painful erection that lasts longer than six hours, as explained by the NIH. This side effect requires immediate medical ention, as it can cause permanent damage if left untreated.

Allergic reaction: Although unlikely, some people may experience an allergic reaction. Signs include a rash or hives, itching, difficulty breathing, and swelling of the face or mouth. Because it may contain other ingredients besides sildenafil as well as contaminants which may also cause allergic reacti

PACKAGING: 4 sachets of 2 orange and 2 pineapple flavors of 5 g, each packed in a cardboard box with insert.

**STORAGE:** Store in a cool, dry place. Protect from light. Keep all medicines out of reach of children.

Manufactured by: Maxwell Life Science Pvt. Ltd. T-53 & T-81, MIDC, TARAPUR, BOISAR District: Palghar - 401506

Manufactured for: CASTILLO INTERNATIONAL GROUP Registered Office 321777 State of Kuwait, Sharq, Block 5, Ahmad Al Jaber Street, Hamad Tower, Floor 12

قد سبيب حيلي عن طريق الفر ( Seedios (rat Jelly) انخفاضيا مذقبًا طفيفًا في ضغط الدو ستحتاج إلى الخضوع لفحص طبي شامل لتشخيص ضعف الانتصاب لديك ومعرفة ما إذا كان من الأمن تناول جيلي عن طريق الفم ( (Seedios Oral Jelly بمفرده أو مع أدويتك الأخرى

تأكد من إخبار طبيبك إذا كنت: النيت من مشاكل في القلب (على سبيل المثال، الذبحة الصدرية، أو ألم الصدر، أو قصور القلب، أو عدم انتظام ضربات الظب، أو النوبة القلبية، أو تضييق الصمام الأورطي)

عانیت من سکتهٔ دماغیهٔ

ضغط دم منخفض أو مرتفع
عانيت من فقدان شديد للرؤي

→ حروب - حروب
→ تعلق من مرض رو (لقي نائر في العين يسمى النهاب الشبكية الصباغي.
→ علتيت من مشاكل في الكلي
→ علتيت من مشاكل في الكليد

•عانيت من مشاكل في الدم، بما في ذلك فقر الدم المنجلي أو سرطان الدم لديك حساسية من السيلدينافيل (المادة الفعالة)

لديك تشوه في القضيب، أو مرض بيروني، أو كان لديك انتصاب استمر لأكثر من ٤ ساعات

تماني من قرحة في المدة أو أي نرع من مذاكل التريف تتناول أدري بكر من من من من و النصي و النوية أن تؤثر على طريقة عمل جبلي عن طريق الفم ( (Seedios Oral Jelly أخير طبيك عن الأدرية التي تتناول أدرية أنو تتوقف عن تتاول الدواء قبل النشارة طبيك. استشارة طبيك. لا ينبغي أبذًا استخدام جيلى عن طريق الله ( ( Seedios Oral Jelly مع الأدوية التي تحتوي على النترات. إذا كنت تتناول ادوية تسمى حاصر ات الله العاج ارتفاع ضغط الدم أو مشاكل البروستاته فقد ينخفض ضغط دمك فجاة.

قا تشرر الدوار أن الإعمام. إذا كنت تقدّل مقمك البرويقز أو إذا كان من المكن تحزل جر علك. لا ينفي استخار مقبق عن طريق الفر ( Secios Grd Jely ) فلاجات الطبية الأخرى التي تسبب الانتصاب نشمل هذه العلاجات الحروب أو الأدوية التي يتم حقتها أو إنخالها في القضيب أو الغرسات أو مضخات القريع.

جيلي عن طريق للغم ( Seedios Oral Jelly ) مخصص فقط للمرضي الثنين يعانون من ضعف الانتصاب لا يُستخد للمواليد الجد أو الأطفل أو النساء. لا يحميك جيلي عن طريق الغم الأمراض المنفولة جنسباء بما في ذلك فيروس نقص المناعة البشرية - القروس السبب لمرض الإيدز. فهو ليس مرمونًا أو مثورا للثموة الجنسية. ( Seedios Oral Jelly أنت أو شريكك من

لحمل والرضاعة الطبيعية والخصبوبة

العصفي الرضعة الميتية وتحصور على الفر ( Sectios Oral Jelly للنساء. لا توجد در اسك كالية وخاصعة للرقابة بشكل جيد على النساء الحوامل أو المرضعك. لم يتم العُور على أي النار جلنبية ذات صلة في در اسك التكثار في الفنران والارانب بعد تناول السولديناقيل عن طريق الفم.

الإنار الجانبية. الإنار الجانبية الأكثر شوغا: في التجارب السريرية التي **منت الموافقة عليها مسيقاً باستخدام** السليدانيال. كما هر مفصل في Model كن التأثير الجانبي الأكثر شوغا هو الصداع، والذي عاني منه ما يصل إلى ٢١٪ من المشاركين في بعد الدرسان، تشمل الأثار الجانبية الأخرى الملغ حقيا في التجارب السريرية قبل الموافقة والتصمن التي حشى بدا اللوفقة الدوغة واحمرار الوجه وعسر الهضم واضطراب المدة والإسهال واحقان الأض والقباب الحلق. مشكل القلب والأرعي الدموية، يمكن أن سبب السيلايتاقيل مشكل للأشخاص المصليين بلمراض القلب والأو مية العربية، ولن أن سبب السيلايتاقيل مشكل القلب والأوعية.

مرجع سطح المكتب (PDR). لذا ارتبط بنبضات قلب سريعة أو غير منتظمة وبالتالي قد لا يكون أمنًا للأشخاص المصابين بأمراض القلب والذبحة الصنرية وانخفاض أو ارتفاع ضغط الدم وعدم انتظام ضريات القلب أو أي شخص عاتى من نوبة قلبية و سكنة دماغية في الأشهر السنة الماضية.

ار سكة داغية في الاثير السلة السنيه. على بعض الرجل، على يعمن الرجل، وفن الصر المقامي بعد تنارل السليديقل، وفقا جي اس الماهد الوطنية المحمة (NHH). يتشل الاثر الجانيية الصيرة الأخرى التى أيغ عنها ستخمر السلية المائة في معنى العالات. القالي: في ريطانية المحررة الأخرى التى أيغ عنها ستخمر السلية الروية، والعين الأكل حسابية السوء ، والتقراب في روية الأوان ، مثل روية لين زرق على الأشاء او صعرية التعييز بين الأخصر والأرزق. تشل الأثر الجانيية الصيرة الأخرى التى أيغ عنها ستخمر السلية العائم في معنى العالات. القالي: في روية الأخرى الذي الذي على المركز من المائة المعالية المحمد والعربي الأكل من ساعات ، كما والصيرة التقراب العالم الان الحقوم على التعليم مؤرم المائع التي التعليم مؤرم المعارب والدينة، والعين الأكل والعنوان العام الاتي الجاني على قرية العيق وسليم على الع القالي: فريز المعالية المركزات التي ولنع المعالية العمر المعاون الارية، والمحمد المعامي العالم العملية العمر على جلب السلينية إلى المركزات التي إلا تعمل الحسية.

12-1-11

لتعبنة والتغليف: ٤ أكياس من نكهتين من البرتقال ونكهتين من الأناناس بوزن ٥ جرام، كل منها معبأة في صندوق من الورق المقوى مع ملحق. التخزين: قم بتخزينه في مكان بار د وجاف احم احفظ جميع الأدوية بعيدًا عن متناول الأطفال ف احمه من الضو ء

	بمصنعة.
العالمية للتجارة العامة والمقاولات	لشركة كاستلو
TTIVVI	مكتب مسجل ا
شرق ، قطعة ٥.	دولة الكويت ،
بابر ، برج حمد ، الدور ١٢	شآرع أحمد الم

ماكسويل لايف ساينس بر ايفت المحدودة. (Maxwell Life Science Pvt. Ltd)

تى-٥٣ وتى-٨١ ميدك ، تار ابور . بواسار ، مقاطعة بالغار -506 401